

*Isolation and characterization of gut  
bacteria of fruit fly, Bactrocera tau  
(Walker)*

**Chandra S. Prabhakar, Pankaj Sood,  
Sarbjit S. Kanwar, Prem N. Sharma, Ajay  
Kumar & Pawan K. Mehta**

**Phytoparasitica**


ISSN 0334-2123

Volume 41

Number 2

Phytoparasitica (2013) 41:193-201

DOI 10.1007/s12600-012-0278-5



**Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media Dordrecht. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.**

# Isolation and characterization of gut bacteria of fruit fly, *Bactrocera tau* (Walker)

Chandra S. Prabhakar · Pankaj Sood · Sarbjit S. Kanwar ·  
 Prem N. Sharma · Ajay Kumar · Pawan K. Mehta

Received: 13 February 2012 / Accepted: 29 November 2012 / Published online: 22 December 2012  
 © Springer Science+Business Media Dordrecht 2012

**Abstract** Gut bacteria of fruit fly, *Bactrocera tau* (Walker) (Diptera: Tephritidae), were isolated and the isolates attractive to *B. tau* adults were characterized using morphological, biochemical and 16S rRNA

analyses to determine their taxonomic position. Based upon morphological, biochemical and 16S rRNA sequences (on the basis of closest match), five gut bacterial species of *B. tau* were characterized as *Delftia acidovorans*, *Pseudomonas putida*, *Flavobacterium* sp., *DeFluvibacter* sp. and *Ochrobactrum* sp., of which four bacterial isolates, viz., *Delftia acidovorans*, *Flavobacterium* sp., *DeFluvibacter* sp. and *Ochrobactrum* sp. are new records from guts of the fruit fly species.

C. S. Prabhakar (✉) · P. Sood · P. K. Mehta  
 Department of Entomology,  
 CSK Himachal Pradesh Krishi Vishvavidyalaya,  
 Palampur, Kangra,  
 Himachal Pradesh 176 062, India  
 e-mail: csprabhakar.ento@gmail.com

P. Sood  
 e-mail: pankajplp@rediffmail.com

S. S. Kanwar · A. Kumar  
 Department of Microbiology,  
 CSK Himachal Pradesh Krishi Vishvavidyalaya,  
 Palampur, Kangra,  
 Himachal Pradesh 176 062, India

P. N. Sharma  
 Molecular Plant Pathology Laboratory, Department of Plant  
 Pathology, CSK Himachal Pradesh Krishi Vishvavidyalaya,  
 Palampur, Kangra,  
 Himachal Pradesh 176 062, India

C. S. Prabhakar  
 International Crops Research Institute for the Semi Arid  
 Tropics, Patancheru,  
 Andhra Pradesh 502 324, India

P. Sood  
 Crop Research Sub Station,  
 CSK Himachal Pradesh Krishi Vishvavidyalaya,  
 Sundernagar, Mandi,  
 Himachal Pradesh 175 019, India

**Keywords** *DeFluvibacter* · *Delftia* · *Flavobacterium* ·  
*Ochrobactrum* · Phylogeny · *Pseudomonas* · 16S rRNA

## Introduction

The insect-bacterial association has co-evolved for more than 250 million years and has resulted in multifaceted interactions between insects and bacteria, ranging from pathogenicity to highly sophisticated mutualistic relationships (Douglas & Beard 1997; Oliver *et al.* 2005; Wernegreen 2002); may be extracellular or intracellular and play a role in the nutrition, physiology and reproduction of the host insect (Brune 1998; Douglas *et al.* 2001). On insect gut bacterial associations, the earliest report was published by Petri in 1909 in an insect species, the olive fly, *Bactrocera* (*Dacus*) *oleae* (Rossi).

A range of bacteria belonging to different genera, viz. *Acetobacter*, *Agrobacterium*, *Arthrobacter*,

*Bacillus*, *Citrobacter*, *Enterobacter*, *Erwinia*, *Escherichia*, *Hafnia*, *Klebsiella*, *Kluyvera*, *Lactobacillus*, *Listeria*, *Micrococcus*, *Pantoea*, *Pectobacterium*, *Proteus*, *Providencia*, *Pseudomonas*, *Raoultella*, *Serratia*, *Staphylococcus*, *Streptococcus* and *Xanthomonas*, has been isolated and characterized from the gut of tephritid fruit fly species (Behar *et al.* 2009; Lloyd *et al.* 1986; Prabhakar *et al.* 2009a; Wang *et al.* 2011).

In fruit flies, symbionts act as a natural source of nitrogen, amino acids and vitamins lacking in fruit tissues to their host and are vertically transmitted to the next generation (Drew *et al.* 1983; Gupta & Anand 2003). Moreover, cultivable gut bacteria have several implications in pest management strategies, *e.g.* bacteria were found to be involved in the degradation of the toxic substances ingested by the host insect leading to insecticide resistance (Bousch & Matsumara 1967; Prabhakar *et al.* 2008). Certain components of bacterial odor play a vital role in fruit fly behavior as either feeding or ovipositional stimulants (Lauzon *et al.* 2000) and are being exploited in pest management in the form of baits or traps (Robacker 2007; Sood *et al.* 2010; Thaochan & Chinajariyawong 2011).

The pumpkin fly, *Bactrocera tau* (Walker), infests a wide range of commercially important cucurbit and solanaceous crops and has a wide distribution throughout south Asian countries (India, Sri Lanka, Bangladesh and Bhutan) to the southeast Asian countries (Thailand, Malaysia, Vietnam, Philippines and Indonesia) and the far east Asian region including Taiwan and south China (Prabhakar *et al.* 2009b, 2012; Thanaphum & Thaenkham 2003; White & Elson-Harris 1992). In the previous study, members of the Enterobacteriaceae (*Klebsiella oxytoca* and *Pantoea agglomerans*) were found to be the dominant microbial population in the gut of laboratory reared *B. tau* (Prabhakar *et al.* 2009a). Laboratory studies on bacterial odors (*K. oxytoca* and *P. agglomerans*) attracting *B. tau* adults (males and females) suggested their possible applied role in fruit fly management programmes (Sood *et al.* 2010). However, there is still a dearth of knowledge on the bacterial communities attracting *B. tau*. The cultivable gut bacteria in particular will be more useful than non-cultivable bacteria when searching for a good fruit fly attractant. Therefore, in the present study, an attempt has been made to characterize cultivable gut bacteria of *B. tau*, a serious pest of cucurbits in southeast Asia, India and China.

## Materials and methods

**Isolation of gut bacteria** The bacteria were isolated from 10-day-old adult flies reared from field-collected cucurbit fruits from vegetable fields of Himachal Pradesh (Table 1), as described by Lloyd *et al.* (1986). Flies were cold anaesthetized for 5 min. and surface sterilized with alcohol (70%) for 30 s. followed by sodium hypochloride (0.25%) for 1 min. and then washed three times with sterilized distilled water (SDW) to remove external contaminations. The surface sterilized flies ( $n=5$  adult flies from each *B. tau* isolate collected from the same location and host plant;  $N=45$  adult flies from nine *B. tau* isolates) were dissected open in physiological saline to remove fly gut. Gut content was streaked separately on Peptone Yeast Extract Agar (PYEA) and Brain Heart Infusion Agar (BHIA) for bacterial growth at  $30\pm1^\circ\text{C}$  for 48–72 h. A single colony of each isolate was used for establishment of pure culture with respective medium and then maintained on PYEA slants and/or plates at 4–8°C.

**Selection of gut bacteria for characterization** Initially, the isolated bacteria were evaluated for their attractancy to fruit fly, *B. tau* (choice method). Thirty gut bacterial isolates, which were frequently isolated from different fruit fly samples across the locations, were used to study their attractiveness to *B. tau*. Pure culture (72 h old) of different bacterial isolates was grown on PYE (peptone yeast extract) broth medium. The bacterial isolates were taken in separate petri plates and kept inside the cage (45 cm×45 cm×55 cm) with un-inoculated PYE broth as control. Twenty-five pairs of 5-day-old fruit flies were released in the cage and flies visiting each treatment were recorded for 30 min. The experiment was repeated six times for *B. tau* and data obtained were analyzed statistically by using the technique of analysis of variance for completely randomized design as described by Gomez & Gomez (1984). The treatment means were compared at 5% level of significance by least significance difference test described by Gomez & Gomez (1984). Five gut bacteria having maximum fruit fly attractiveness were selected for further characterization.

## Characterization of gut bacteria

**Morphological and biochemical characterization** Morphological (shape, Gram's staining, mobility), cultural

**Table 1** Isolation of gut bacteria from different populations of *Bactrocera tau* from Himachal Pradesh (India)

| <i>B. tau</i> isolates ( <i>n</i> =5) used for bacterial isolation |                      |               | Culture media used                |                                                       |                          |                                  |                                                       |                          |
|--------------------------------------------------------------------|----------------------|---------------|-----------------------------------|-------------------------------------------------------|--------------------------|----------------------------------|-------------------------------------------------------|--------------------------|
| Isolate No. of <i>B. tau</i>                                       | Location (District)  | Host          | Peptone Yeast Extract Agar (PYEA) |                                                       |                          | Brain Heart Infusion Agar (BHIA) |                                                       |                          |
|                                                                    |                      |               | Bacterial colonies isolated       | Bacterial isolates selected for attractancy screening | Bacterial isolate number | Bacterial colonies isolated      | Bacterial isolates selected for attractancy screening | Bacterial isolate number |
| P1                                                                 | Nadaun (Hamirpur)    | Bottle gourd  | 4                                 | 4                                                     | P1A, P1B, P1C, P1D       | 4                                | 2                                                     | B1A, B1B                 |
| P2                                                                 | Nadaun (Hamirpur)    | Bitter gourd  | 3                                 | 2                                                     | P2A, P2B                 | 4                                | 1                                                     | B2A                      |
| P3                                                                 | Sundernagar (Mandi)  | Cucumber      | 3                                 | 1                                                     | P3A                      | 3                                | 2                                                     | B3A, B3B                 |
| P4                                                                 | Palampur (Kangra)    | Cucumber      | 2                                 | 1                                                     | P4A                      | 5                                | 2                                                     | B4A, B4B                 |
| P5                                                                 | Nauni (Solan)        | Summer squash | 4                                 | 2                                                     | P5A, P5B                 | 4                                | 1                                                     | B5A                      |
| P9                                                                 | Mandi (Mandi)        | Cucumber      | 2                                 | 1                                                     | P9A                      | 3                                | 1                                                     | B9A                      |
| P10                                                                | Ghumarwin (Bilaspur) | Cucumber      | 4                                 | 2                                                     | P10A, P10B               | 4                                | 2                                                     | B10A, B10B               |
| P15                                                                | Jawalamukhi (Kangra) | Bitter gourd  | 2                                 | 1                                                     | P15A                     | 5                                | 2                                                     | B15A, B15B               |
| P18                                                                | Shahpur (Kangra)     | Bitter gourd  | 4                                 | 2                                                     | P18A, P18B               | 3                                | 1                                                     | B18A                     |
| Total                                                              |                      |               | 28                                | 16                                                    |                          | 35                               | 14                                                    |                          |

(pigment production, growth in broth medium) and biochemical (citrate, methyl red, VP, TSI, catalase, oxidase and carbohydrate fermentation tests) characterization of pure culture was done by standard techniques and isolates were identified using *Bergey's Manual of Determinative Bacteriology* (Holt *et al.* 2000).

#### Molecular characterization

**Extraction of genomic DNA** Total genomic DNA of each isolate was extracted following the method of Prabhakar *et al.* (2009a). The 48-h-old bacterial culture multiplied on Peptone Yeast Extract Broth (PYEB) was transferred to 1.5 ml microtubes and spun at 10,000 rpm for 12 min. After discarding the supernatant, the microtubes containing bacterial pellets (approx. 50 mg) were immersed in liquid nitrogen container for one min and the pellet was ground to fine powder immediately using a micro pestle. To this, 700  $\mu$ l of cetyltrimethyl ammonium bromide (CTAB) extraction buffer was added and incubated at 65°C for 1 h in a water bath (York Scientific Industries, Delhi, India).

An equal volume (700  $\mu$ l) of chloroform: isoamyl alcohol (24:1 v/v) was added and the contents were mixed thoroughly. Tubes were spun at 10,000 rpm for 12 min in a high speed refrigerated centrifuge (Remi Elektrotechnik, India) at 4°C. Aqueous phase was transferred to new tubes and 450  $\mu$ l pre-chilled iso-propanol was added and kept at –20°C for 20–30 min to precipitate the DNA. Tubes were then spun at 10,000 rpm for 12 min and supernatant was decanted. The DNA pellet was washed with 70% ethanol (three times), dried and dissolved in 100  $\mu$ l of Tris EDTA buffer (10 mM Tris HCl and 1 mM EDTA, pH 8.0). RNase @ 10  $\mu$ lml<sup>–1</sup> (MBI Fermentas) was added and the emulsion was incubated for half an hour at 37°C. The amount of DNA was quantified by recording the absorbance at 260 nm wavelength using an UV/VIS spectrophotometer (Bio Rad, SmartSpec 3000). DNA was stored at –20°C for further use.

**PCR amplification and sequencing** Polymerase chain reaction (PCR) was performed with eubacterial primers pair forward 5'AGAGTTTGATCATGGCTCAG3' and reverse 5' TACCTTGTTACGACTTCACC 3' targeting



the 1,400 bp *rrs* (16S rRNA) gene (Heddi *et al.* 1998). The PCR amplification was carried out in 0.2 ml PCR tubes with 25 µl reaction volume containing 10 ng of DNA template, 20 pmol of each primer in 25 mM MgCl<sub>2</sub>, 10 mM of each deoxyribonucleoside triphosphate (Fermentas), 5 units of *Taq* polymerase (Life Technologies India, Pvt. Ltd.) and 10X reaction buffer. Amplifications were performed using a thermal cycler (GeneAmp PCR system 9700, Applied Biosystems, Carlsbad, CA, USA) with an initial denaturation step of 5 min at 94°C followed by 35 cycles at 94°C for 45 s, 53°C for 45 s, 72°C for 30 s and a final elongation step at 72°C for 5 min. The product was separated in a 1% (w/v) agarose gel in TAE buffer (40 mM Tris-acetate, 1 mM EDTA). PCR products (1,450 bp) of 16S rRNA gene of five gut bacteria obtained through amplification with specific primers were first purified with ethanol precipitation of amplified DNA, freeze dried (Christ *Alpha I-2LD*), and then custom sequenced (ABI PRISM 310<sup>TM</sup> Genetic Analyzer, Applied Biosystems, USA) using the same upstream and downstream primers (Life Technologies India).

**Nucleotide sequence analysis** The sequences of different bacterial isolates were blasted using online NCBI Blastn program (<http://www.ncbi.nih.gov/blast>). For the purpose, 42 sequences of 16S rRNA of different bacteria of high sequence similarity were selected for sequence comparison from GenBank Nucleotide Database, NCBI. The pair-wise genetic distance between five gut bacterial isolates of *B. tau* and other selected bacterial sequences was determined.

The evolutionary history was inferred using the UPGMA method (Sneath & Sokal 1973). The evolutionary distances were calculated using Maximum Composite Likelihood method with bootstrap test (500 replicates) (Tamura *et al.* 2004). All positions containing gaps and missing data were eliminated from the dataset (complete deletion option). There were a total of 298 positions in the final dataset. Phylogenetic analysis was conducted in MEGA 4.1 Software programme (Tamura *et al.* 2007).

## Results

**Isolation and identification of gut bacterial isolates of fruit fly** Gut bacteria were isolated from nine populations of fruit fly, *B. tau* on two culture media, viz.

PYEA and BHIA (enriched culture media). It was found that bacteria were associated with all the nine populations of *B. tau* (Table 1). A total of 63 different bacterial colonies were observed on two culture media. Of these, 28 were isolated on PYEA and 35 on BHIA. On the basis of colony morphology (pigmentation, shape and size of colony), 16 and 14 isolates obtained on PYEA and BHIA (which were most common among different fruit fly isolates across the locations), respectively, were evaluated for the attractancy to adult fruit flies under laboratory conditions to select the most promising gut bacteria in terms of their attractancy for further characterization.

The five most promising gut bacterial isolates were selected based on their attractancy to *B. tau* (Table 2) for further characterization. These five gut bacterial isolates: P1B, P3A, P10A, B4A and B10B (P/B-media used for isolation, i.e., PYEA and Brain Heart Infusion Agar; 1,3,10,4.10: *B. tau* isolate numbers giving information about the location and host; and A,B,C,...: different bacterial colonies isolated from the gut of *B. tau*) were gram negative and rod shaped. All five isolates of gut bacteria were producing sediments when multiplied in broth medium. All the five gut bacterial isolates were non-pigment producing and showed mobility except for P10A, which produced yellow pigment on PYEA plates and was non-mobile. With regard to biochemical characteristics, all the isolates showed negative reaction for citrate, VP, TSI and gas production in glucose medium tests, whereas they were found positive for catalase and oxidase test. All isolates except PIB were negative for methyl red test (Table 3). On the basis of cultural, morphological, biochemical and 16S rRNA gene characteristics, the closest match to isolate P1B, P3A, P10A, B4A and B10B was *Delftia acidovorans* (91% DNA identity), *Pseudomonas putida* (97% DNA identity), *Flavobacterium* sp. (95% DNA identity), *Defluviobacter* sp. (98% DNA identity) and *Ochrobactrum* sp. (99% DNA identity), respectively (Table 3).

The 16S rRNA gene nucleotide sequences of these isolates were submitted to GenBank nucleotide database under accession numbers HQ446523 to HQ446527.

**Phylogenetic studies of gut bacteria** All the sequences of five bacterial isolates were compared with 42 other bacterial sequences available online in GenBank (NCBI) by multiple sequence alignment tools using

**Table 2** Attractancy of bacterial isolates against fruit fly, *Bactrocera tau*

| Sr. No. | Bacterial isolates (72 h old, 2 ml broth culture) | Fruit flies visited/30 min (n=50) |       |        |
|---------|---------------------------------------------------|-----------------------------------|-------|--------|
|         |                                                   | Female*                           | Male* | Total* |
| 1       | P1A                                               | 3.17                              | 2.17  | 5.33   |
| 2       | P1B**                                             | 9.83                              | 7.17  | 17.00  |
| 3       | P1C                                               | 3.50                              | 3.83  | 7.33   |
| 4       | P1D                                               | 3.17                              | 3.50  | 7.17   |
| 5       | P2A                                               | 1.17                              | 1.00  | 2.17   |
| 6       | P2B                                               | 3.00                              | 2.00  | 5.00   |
| 7       | P3A**                                             | 10.5                              | 7.67  | 18.17  |
| 8       | P4A                                               | 3.50                              | 1.33  | 4.83   |
| 9       | P5A                                               | 4.33                              | 3.00  | 7.33   |
| 10      | P5B                                               | 3.83                              | 1.83  | 5.67   |
| 11      | P9A                                               | 5.17                              | 4.50  | 10.50  |
| 12      | P10A**                                            | 8.67                              | 6.67  | 15.33  |
| 13      | P10B                                              | 4.67                              | 4.83  | 9.50   |
| 14      | P15A                                              | 2.17                              | 2.50  | 4.67   |
| 15      | P18A                                              | 1.67                              | 1.33  | 2.17   |
| 16      | P18B                                              | 3.50                              | 4.17  | 7.67   |
| 17      | B1A                                               | 2.33                              | 3.83  | 6.17   |
| 18      | B1B                                               | 3.00                              | 2.50  | 5.17   |
| 19      | B2A                                               | 3.50                              | 3.17  | 9.50   |
| 20      | B3A                                               | 3.33                              | 2.67  | 6.00   |
| 21      | B3B                                               | 2.17                              | 2.50  | 4.67   |
| 22      | B4A**                                             | 7.50                              | 6.33  | 13.83  |
| 23      | B4B                                               | 2.67                              | 2.83  | 5.50   |
| 24      | B5A                                               | 3.83                              | 1.50  | 5.33   |
| 25      | B9A                                               | 2.50                              | 3.00  | 5.50   |
| 26      | B10A                                              | 2.50                              | 1.33  | 4.00   |
| 27      | B10B**                                            | 6.33                              | 5.83  | 12.17  |
| 28      | B15A                                              | 4.33                              | 4.17  | 8.83   |
| 29      | B15B                                              | 3.00                              | 2.83  | 6.17   |
| 30      | B18A                                              | 2.67                              | 2.17  | 4.83   |
| 31      | Control (Un-inoculated PYEA broth)                | 1.67                              | 1.83  | 3.50   |
|         | LSD <sub>0.05</sub>                               | 1.58                              | 1.48  | 2.37   |

\*Mean of six replications, LSD test

\*\*Selected for characterization

ClustalW programme. The per cent pair-wise genetic distance of the five promising isolates with other bacterial sequences ranged from 0.00 to 0.70 nucleotide per site.

The dendrogram constructed by phylogenetic analysis presented in Fig. 1 showed that all the bacterial isolates, viz., P1B, P3A, B4A and B10B, clustered with *Delftia*, *Pseudomonas*, *Deftuvibacter* and *Ochrobactrum*, respectively (all Proteobacteria), except P10A (HQ446525), which was clustered with *Flavobacterium* (a typical Bacteroidetes). Based on their affinity with known sequences in databank, the isolate P1B belongs to class  $\beta$ -Proteobacteria, P3A to class  $\gamma$ -Proteobacteria, and B4A & B10B to class  $\alpha$ -Proteobacteria.

## Discussion

Many workers suggested that nucleic acid sequence approaches, particularly 16S rRNA genes, have proved an important tool to settle the taxonomic position of the microbial community of insects (Brauman *et al.* 2001; Wang *et al.* 2011). An immense library of sequence data for 16S rRNA loci and other robust markers, allows the precise identification of many associated species, even those that resist cultivation (Stevenson *et al.* 2004). Over 200,000 bacterial entries exist currently for 16S rRNA, and these sequences can place most surveyed bacterial taxa securely into genera, if not species (Rupp 2004; Ueda *et al.* 2004).

The gut of tephritid fruit flies in general (Behar *et al.* 2009; Lloyd *et al.* 1986; Wang *et al.* 2011) and *B. tau* in particular (Prabhakar *et al.* 2009a; Sood & Nath 2002) is a storehouse of bacterial community. In the present study also 30 different bacterial colonies on the basis of colony morphology were screened for attractancy, but only five isolates which were found promising in terms of their attractancy to adult *B. tau* flies and could have a great implication on their future management programmes were further characterized. All these bacterial isolates except *P. putida* were not reported earlier to be associated with *B. tau* and having any role in fruit fly ecology. The attractiveness of these isolates to fruit flies, however, suggests their possible role in the fruit fly nutrition and physiology. Concrete and concentrated efforts across the fruit fly species are needed to elucidate the complex phenomenon to some conclusion.

*Delftia* was isolated from *B. tau* for the first time, although its association was earlier reported with cotton bollworm, *Helicoverpa armigera* (Hübner) (Xiang *et al.* 2006); wood borer, *Saperda vestita* (Say) (Delalibera

**Table 3** Morphological, biochemical and molecular characteristics of gut bacteria of *Bactrocera tau*

| Characteristics                    | Bacterial isolates                  |                                    |                                    |                                    |                                  |
|------------------------------------|-------------------------------------|------------------------------------|------------------------------------|------------------------------------|----------------------------------|
|                                    | P1B                                 | P3A                                | P10A                               | B4A                                | B10B                             |
| <b>Morphological</b>               |                                     |                                    |                                    |                                    |                                  |
| Shape                              | Rod                                 | Rod                                | Rod                                | Rod                                | Rod                              |
| Gram's reaction                    | –                                   | –                                  | –                                  | –                                  | –                                |
| Pigment production                 | –                                   | –                                  | Y                                  | –                                  | –                                |
| growth in broth medium             | Sediment                            | Sediment                           | Sediment                           | Sediment                           | Sediment                         |
| Mobility                           | +                                   | +                                  | –                                  | +                                  | +                                |
| <b>Biochemical</b>                 |                                     |                                    |                                    |                                    |                                  |
| Citrate test                       | –                                   | d                                  | –                                  | –                                  | d                                |
| Methyl red                         | +                                   | –                                  | –                                  | –                                  | –                                |
| V P test                           | –                                   | d                                  | –                                  | –                                  | –                                |
| TSI                                | –                                   | –                                  | –                                  | –                                  | –                                |
| Catalase                           | +                                   | +                                  | +                                  | +                                  | +                                |
| Oxidase                            | +                                   | +                                  | +                                  | +                                  | +                                |
| D-Glucose                          | –                                   | +                                  | +                                  | –                                  | d                                |
| Gas production in glucose medium   | –                                   | –                                  | –                                  | –                                  | –                                |
| <b>Molecular</b>                   |                                     |                                    |                                    |                                    |                                  |
| 16S rDNA sequence blast similarity | 91% with <i>Delftia acidovorans</i> | 97% with <i>Pseudomonas putida</i> | 95% with <i>Flavobacterium</i> sp. | 98% with <i>Deffluvibacter</i> sp. | 99% with <i>Ochrobactrum</i> sp. |
| Bacteria–closest match to          | <i>Delftia acidovorans</i>          | <i>Pseudomonas putida</i>          | <i>Flavobacterium</i> sp.          | <i>Deffluvibacter</i> sp.          | <i>Ochrobactrum</i> sp.          |

Y yellow pigment, – negative reaction, + positive reaction, d doubtful result


*et al.* 2005); tobacco caterpillar, *Manduca sexta* (Brinkmann *et al.* 2008); and mosquito, *Aedes albopictus* Skuse (Zouache *et al.* 2009). Whereas *Pseudomonas putida* and members of genera *Pseudomonas* have been reported from many insect species including fruit flies as well as from *B. tau* (Brinkmann *et al.* 2008; Sood & Nath 2002). The presence of *Flavobacterium* sp., *Deffluvibacter* sp. and *Ochrobactrum* sp. has also been reported in other insect species, viz. *Flavobacterium* sp. from ant, *Tetraponera binghami* Forel (van Borm *et al.* 2002), honey bees (Mohr & Tebbe 2007) and tobacco caterpillar, *M. sexta* (Brinkmann *et al.* 2008); however, the present findings constitute the first report of their association with fruit flies, particularly *Bactrocera* species.

*Deffluvibacter* sp. is a member of the bacterial family Phyllobacteriaceae from the class  $\alpha$ -Proteobacteria and has not been reported from the gut of any insect species. An unassigned bacterium (member Phyllobacteriaceae) has, however, been reported from the

gut content of Asian longhorned beetle, *Anoplophora glabripennis* Motschulsky (Geib *et al.* 2009). The bacterial family Phyllobacteriaceae is closely related to the Bradyrhizobiaceae, Methylobacteriaceae and Rhizobiaceae. Bacteria from the families Methylobacteriaceae and Rhizobiaceae have been reported from different insect species, viz. *Rhizobium* and *Methylobacterium* from the gut of *Tetraponera* ants (van Borm *et al.* 2002) and *Rhizobium* from the gut content of Asian longhorned beetle, *Anoplophora glabripennis* (Geib *et al.* 2009).

*Ochrobactrum* sp. belongs to the  $\alpha$ -2 subclass of the Proteobacteria (De Ley 1992). The genus was first described by Holmes *et al.* (1988) and the phylogenetic position of *Ochrobactrum* sp. was defined by De Ley (1992) and Yanagi & Yamasato (1993) on the basis of DNA  $\pm$  rRNA hybridization and 16S rRNA homology studies. Its closest known relative is *Brucella* (De Ley 1992; Yanagi & Yamasato 1993).





**Fig. 1** Phylogenetic tree based on 16S rRNA gene sequences using Unpaired Group Mean Algorithm (UPGMA) method showing the relationships between five gut bacterial isolates of *Bactrocera tau*.

Numbers above the nodes are bootstrap values (500 replicates). Phylogenetic analyses were conducted in software MEGA 4.1. Arrows indicate gut bacteria of *B. tau* isolated in present study

*Ochrobactrum* sp. was reported from the insect gut (Asian longhorned beetle, *Anoplophora glabripennis*) by Geib *et al.* (2009); its closest relative, *Brucella* sp., was isolated from the gut of wood borer *Saperda vestita*

and identified by 16S rRNA analysis by Delalibera *et al.* (2005).

Earlier *Klebsiella oxytoca*, *Pantoea agglomerans*, *Staphylococcus* sp., *Pseudomonas putida*, *Bacillus* sp.

and *Enterobacter agglomerans* were reported from the gut of fruit fly, *B. tau* (Prabhakar *et al.* 2009a; Sood & Nath 2002). The present finding added a few more genera to this list, but as a whole all these reports suggested that a vast range of gut bacterial diversity exists in the *B. tau* system. Thorough isolation and characterization with culture-dependent and -independent techniques are therefore needed to explore the gut bacterial diversity of *B. tau* to understand the host behavior in relation to gut bacterial community. On the other hand, vertebrate pathogenicity of these bacteria can also not be denied as some species of *Ochrobactrum*, *Delftia*, *Staphylococcus* and *Deftuvibacter* were previously reported as human pathogens (Battaglia 2008; Mastroianni *et al.* 1999; Ogston 1984; Preiswerk *et al.* 2011); this, however, requires further investigations. Detailed investigations are also needed to establish the taxonomic positions of *Delftia*, *Flavobacterium* sp., *Deftuvibacter* sp. and *Ochrobactrum* sp. up to species level using chemotaxonomic and molecular approaches to understand the tri-trophic interaction among host–fruit flies–microbes.

**Acknowledgments** The authors are grateful to three anonymous reviewers for their helpful critical comments on the manuscript. They also thank Miss Kajal (Senior Research Fellow, Molecular Plant Pathology Laboratory, CSKHPKV, Palampur) for the help rendered during the study.

## References

- Battaglia, T. C. (2008). *Ochrobactrum anthropi* septic arthritis of the acromioclavicular joint in an immunocompetent 17 year old. *Orthopedics*, 31, 606.
- Behar, A., Ben-Yosef, M., Lauzon, C. R., Yuval, B., & Jurkevich, E. (2009). Structure and function of the bacterial community associated with the Mediterranean fruit fly. In K. Bourtzis & T. Miller (Eds.), *Insect symbiosis* (pp. 251–271). Boca Raton, FL: CRC.
- Bousch, G. M., & Matsumura, F. (1967). Insecticidal degradation by *Pseudomonas melophthora*, the bacterial symbiote of the apple maggot. *Journal of Economic Entomology*, 69, 918–920.
- Brauman, A., Dore, J., Eggleton, P., Bignell, D., Breznak, J. A., & Kane, M. D. (2001). Molecular phylogenetic profiling of prokaryotic communities in guts of termites with different feeding habits. *FEMS Microbiology Ecology*, 35, 27–36.
- Brinkmann, N., Martens, R., & Tebbe, C. C. (2008). Origin and diversity of metabolically active gut bacteria from laboratory-bred larvae of *Manduca sexta* (Sphingidae: Lepidoptera: Insecta). *Applied and Environmental Microbiology*, 74, 7189–7196.
- Brune, A. (1998). Termite guts: the world's smallest bioreactors. *Trends in Biotechnology*, 16, 16–21.
- De Ley, J. (1992). The proteobacteria: ribosomal RNA cistron similarities and bacterial taxonomy. In A. Balows *et al.* (Eds.), *The prokaryotes* (pp. 2111–2140). New York, NY: Springer.
- Delalibera, I., Handelsman, J. O., & Raffa, K. F. (2005). Contrasts in cellulolytic activities of gut microorganisms between the wood borer, *Saperda vestita* (Coleoptera: Cerambycidae), and the bark beetles, *Ips pini* and *Dendroctonus frontalis* (Coleoptera: Curculionidae). *Environmental Entomology*, 34, 541–547.
- Douglas, A. E., & Beard, C. B. (1997). Microbial symbiosis in the midgut of insects. In M. Lehane (Ed.), *Biology of the insect midgut* (pp. 315–333). New York: Academic.
- Douglas, A. E., Minto, L. B., & Wilkinson, T. L. (2001). Quantifying nutrient production by the microbial symbionts in an aphid. *Journal of Experimental Biology*, 204, 349–358.
- Drew, R. A. I., Courtice, A. C., & Teakle, D. S. (1983). Bacteria as a natural source of food for fruit flies (Diptera: Tephritidae). *Oecologia*, 60, 279–284.
- Geib, S. M., Jimenez-Gasco, M. D. M., Carlson, J. E., Tien, M., & Hoover, K. (2009). Effect of host tree species on cellulase activity and bacterial community composition in the gut of larval Asian longhorned beetle. *Environmental Entomology*, 38, 686–699.
- Gomez, K. A., & Gomez, A. A. (1984). *Statistical procedures for agricultural research* (2nd ed.). New York: John Wiley & Sons.
- Gupta, K., & Anand, M. (2003). *In vitro* vitamin synthesis by the symbionts of *Bactrocera cucurbitae*. *Indian Journal of Entomology*, 65, 518–528.
- Heddi, A., Charles, H., Khatchadourian, C., Bonnot, G., & Nardon, P. (1998). Molecular characterization of the principal symbiotic bacteria of the weevil *Sitophilus oryzae*: A peculiar G+C content of an endocytobiotic DNA. *Journal of Molecular Evolution*, 47, 52–61.
- Holmes, B., Popoff, M., Kiredjian, M., & Kersters, K. (1988). *Ochrobactrum anthropi* gen. nov., sp. nov. from human clinical specimens and previously known as group Vd. *International Journal of Systematic Bacteriology*, 38, 406–416.
- Holt, J. G., Krieg, N. R., Sneath, P. H. A., Staley, J. T., & Williams, S. T. (2000). *Bergey's manual of determinative bacteriology* (pp. 175–533). New York, NY: Lippincott Williams and Wilkins.
- Lauzon, C. R., Sjogren, R. E., & Prokopy, R. J. (2000). Enzymatic capabilities of bacteria associated with apple maggot flies, a postulated role in attraction. *Journal of Chemical Ecology*, 26, 953–967.
- Lloyd, A. C., Drew, R. A. I., Teakle, D. S., & Hayward, A. C. (1986). Bacteria associated with some *Dacus* species (Diptera: Tephritidae) and their host fruits in Queensland. *Australian Journal of Biological Science*, 39, 361–368.
- Mastroianni, A., Cancellieri, C., & Montini, G. (1999). *Ochrobactrum anthropi* bacteremia: case report and review of the literature. *Clinical and Microbiological Infections*, 5, 570–573.
- Mohr, K. I., & Tebbe, C. C. (2007). Field study results on the probability and risk of a horizontal gene transfer from

- transgenic herbicide-resistant oilseed rape pollen to gut bacteria of bees. *Applied Microbiology and Biotechnology*, 75, 573–582.
- Ogston, A. (1984). On abscesses: classics in infectious diseases. *Review of Infectious Diseases*, 6, 122–128.
- Oliver, K. M., Moran, N. A., & Hunter, M. S. (2005). Variation in resistance to parasitism in aphids is due to symbionts and not host genotype. *Proceedings of the National Academy of Sciences USA*, 102, 12795–12800. doi:10.1073/pnas.0506131102.
- Petri, L. (1909). Ricerche sopra i batteri intestinali della mosca olearia. *Memorie della Regia Stazione di Patologia Vegetale di Roma*, 1–130.
- Prabhakar, C. S., Sood, P., Kapoor, V., Kanwar, S. S., Mehta, P. K., & Sharma, P. N. (2009a). Molecular and biochemical characterization of three bacterial symbionts of fruit fly, *Bactrocera tau* (Tephritidae: Diptera). *Journal of General and Applied Microbiology*, 55, 213–220.
- Prabhakar, C. S., Sood, P., & Mehta, P. K. (2008). Protein hydrolyzation and pesticide tolerance by gut bacteria of *Bactrocera tau* (Walker). *Pest Management and Economic Zoology*, 16, 123–129.
- Prabhakar, C. S., Sood, P., & Mehta, P. K. (2012). Fruit fly (Diptera: Tephritidae) diversity in cucurbit fields and surrounding forest areas of Himachal Pradesh, a North-Western Himalayan state of India. *Archives of Phytopathology and Plant Protection*, 45, 1210–1217. doi:10.1080/03235408.2012.660612.
- Prabhakar, C. S., Sood, P., Mehta, P. K., & Choudhary, A. (2009b). Distribution and developmental biology of fruit flies infesting cucurbits in north-western Himalaya. *Journal of Insect Science*, 22, 300–308.
- Preiswerk, B., Ullrich, S., Speich, R., Bloemberg, G. V., & Hombach, M. (2011). Human infection with *Delftia tsuruhatensis* isolated from a central venous catheter. *Journal of Medical Microbiology*, 60, 246–248.
- Robacker, D. C. (2007). Chemical ecology of bacteria relationships with fruit flies. Integrated Protection of Olive Crops. *IOBC/WPRS Bulletin*, 30, 9–22.
- Rupp, S. (2004). Proteomics on its way to study host-pathogen interaction in *Candida albicans*. *Current Opinion in Microbiology*, 7, 330–335.
- Sneath, P. H. A., & Sokal, R. R. (1973). *Numerical taxonomy*. San Francisco, CA: Freeman.
- Sood, P., & Nath, A. (2002). Bacteria associated with *Bactrocera* sp. (Diptera: Tephritidae)—Isolation and identification. *Pest Management and Economic Zoology*, 10, 1–9.
- Sood, P., Prabhakar, C. S., & Mehta, P. K. (2010). Eco-friendly management of fruit flies through their gut bacteria. *Journal of Insect Science*, 23, 275–283.
- Stevenson, B. S., Eichorst, S. A., Wertz, J. T., Schmidt, T. M., & Breznak, J. A. (2004). New strategies for cultivation and detection of previously uncultured microbes. *Applied and Environmental Microbiology*, 70, 4748–4755.
- Tamura, K., Dudley, J., Nei, M., & Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular and Biological Evolution*, 24, 1596–1599.
- Tamura, K., Nei, M., & Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences USA*, 101, 11030–11035.
- Thanaphum, S., & Thaeankham, U. (2003). Relationships of forms within the *Bactrocera tau* (Walker) (Diptera: Tephritidae) taxon based on *heat shock protein 70* cognate sequences. *Annals of the Entomological Society of America*, 96, 44–53.
- Thaochan, N., & Chinajariyawong, A. (2011). Attraction of *Bactrocera cucurbitae* and *B. papayae* (Diptera: Tephritidae) to the odor of the bacterium *Enterobacter cloacae*. *The Philippine Agricultural Scientist*, 94, 1–6.
- Ueda, K., Yamashita, A., Ishikawa, J., Shimada, M., Watsuji, T. O., Morimura, K., et al. (2004). Genome sequence of *Symbiobacterium thermophilum*, an uncultivable bacterium that depends on microbial commensalism. *Nucleic Acids Research*, 32, 4937–4944.
- van Borm, S., Buschinger, A., Boomsma, J. J., & Billen, J. (2002). Tetraponera ants have gut symbionts related to nitrogen-fixing root-nodule bacteria. *Proceedings of the Royal Society of London B (Biological Sciences)*, 269, 2023–2027.
- Wang, H., Jin, L., & Zhang, H. (2011). Comparison of the diversity of the bacterial communities in the intestinal tract of adult *Bactrocera dorsalis* from three different populations. *Journal of Applied Microbiology*, 110, 1390–1401.
- Wernegreen, J. J. (2002). Genome evolution in bacterial endosymbionts of insects. *Nature Reviews Genetics*, 3, 850–861.
- White, I. M., & Elson-Harris, M. M. (1992). *Fruit flies of economic significance: their identification and bionomics*. Wallingford, UK: Centre for Agriculture and Biosciences International.
- Xiang, H., Wei, G. F., Jia, S., Huang, J., Miao, X. X., & Zhou, Z. (2006). Microbial communities in the larval midgut of laboratory and field populations of cotton bollworm (*Helicoverpa armigera*). *Canadian Journal of Microbiology*, 52, 1085–1092.
- Yanagi, M., & Yamasato, K. (1993). Phylogenetic analysis of the family *Rhizobiaceae* and related bacteria by sequencing of 16S rRNA gene using PCR and DNA sequencer. *FEMS Microbiology Letters*, 107, 115–120.
- Zouache, K., Voronin, D., Tran-Van, V., Mousson, L., Failloux, A. B., & Mavingui, P. (2009). Persistent *Wolbachia* and cultivable bacterial infection in the reproductive and somatic tissues of the mosquito vector *Aedes albopictus*. *PLoS One*, 4, e6388. doi:10.1371/journal.pone.0006388.