

<https://doi.org/10.33307/entomon.v4i2.437>
 ENTOMON 44(2): 111-116 (2019)
 Article No. ent. 44204



Zootaxa 4550 (2): 268-276
<https://www.mapress.com/j/zt/>
 Copyright © 2019 Magnolia Press

Article

<https://doi.org/10.11646/zootaxa.4550.2.7>
<http://zoobank.org/urn:lsid:zoobank.org:pub:7A21D973-61CF-442B-9E43-55B8743B0789>

ISSN 1175-5326 (print edition)
ZOOTAXA
 ISSN 1175-5334 (online edition)

Description of a new species of *Collocleonymus* Masi (Hymenoptera: Chalcidoidea: Pteromalidae) from India

Ankita Gupta, Namitha George¹ and P. M. Sureshan^{1*}

¹ICAR-National Bureau of Agricultural Insect Resources, Hebbal, Bengaluru 560 024, Karnataka, India; ²Zoological Survey of India, Western Ghat Regional Centre, Kozhikode 673006, Kerala, India. Email: pmsuresh43@gmail.com

ABSTRACT: The first Indian species of the genus *Collocleonymus* Masi viz. *C. indiaensis* sp. nov., is formally described from Bengaluru in southern India. © 2019 Association for Advancement of Entomology

Zootaxa 4388 (3): 425-430
<http://www.mapress.com/j/zt/>
 Copyright © 2018 Magnolia Press

Article

ISSN 1175-5326 (print edition)
ZOOTAXA
 ISSN 1175-5334 (online edition)

<https://doi.org/10.11646/zootaxa.4388.3.8>
<http://zoobank.org/urn:lsid:zoobank.org:pub:115FD523-A9F7-4A2C-8243-990D7A9782FB>

A new species of *Acanthormius* (Braconidae: Lysiterminae) reared as a gregarious parasitoid of psychid caterpillars (Lepidoptera: Psychidae) from India

ANKITA GUPTA^{1,2} & DONALD L. J. QUICKE²

¹ICAR-National Bureau of Agricultural Insect Resources, P. B. No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bengaluru, 560 024, India

²Integrative Ecology Laboratory, Department of Biology, Faculty of Science, Chulalongkorn University, Phayathai Road, Pathumwan, BKK 10330, Thailand

*Corresponding author. E-mail: drankitgupta7@gmail.com

Abstract

Acanthormius indicus Gupta & Quicke sp. nov. is described and illustrated. It is a gregarious larval parasitoid of an unidentified bagworm moth caterpillar (Lepidoptera: Psychidae) from southern India. The biology and host associations of members of subfamily Lysiterminae are also discussed.

Zootaxa 4193 (1): 197-200
<http://www.mapress.com/j/zt/>
 Copyright © 2016 Magnolia Press

Correspondence

ISSN 1175-5326 (print edition)
ZOOTAXA
 ISSN 1175-5334 (online edition)

<https://doi.org/10.11646/zootaxa.4193.1.13>
<http://zoobank.org/urn:lsid:zoobank.org:pub:96784CBB-ACB5-4CF7-B455-72424FD55FA5>

The first record of the genus *Tanaostigma* (Hymenoptera: Tanaostigmatidae) in the Old World, with the description of a new species from India

ANKITA GUPTA¹ & SUNIL JOSHI

¹ICAR-National Bureau of Agricultural Insect Resources, P. B. No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bengaluru, 560 024, India

*Corresponding author. E-mail: drankitgupta7@gmail.com

Tanaostigma Howard (Hymenoptera: Tanaostigmatidae) is recorded for the first time in the fauna of the Old World, with *T. indica* Gupta sp. n. described and illustrated from southern India, reared from *Millettia* (=Pongamia) pinnata (Fabaceae).

Key words: *Millettia* (=Pongamia) pinnata, Fabaceae



Journal of Biological Control, 28(3): 122-125, 2014



Research Article

A new species of genus *Cassidibracon* Quicke (Hymenoptera: Braconidae: Braconinae) with new host record from India

ANKITA GUPTA¹ and V. NAVEEN

¹ICAR National Bureau of Agriculturally Important Insects, Hebbal, Bengaluru - 560 024, Karnataka, India.

*Corresponding author. E-mail: drankitgupta7@gmail.com

ABSTRACT: A new species, *Cassidibracon repens* Gupta sp. n. from Karnataka, India, is described and illustrated along with the host caterpillar and cocoons of another closely allied Indian species. The species was reared from cocoons of a Pyralidae caterpillar which is a new host record for *Cassidibracon*. The main diagnostic characters include - second, third and fourth metasomal tergites with midlongitudinal carina; ovipositor length 0.53× length of gaster; OOL/POL = 2.5; eye length/malar space = 3.33; median flagellomeres approximately 2-2.27× longer than wide and antenna with 24 flagellomeres. Revised Key to the Oriental species of the Genus *Cassidibracon* Quicke is also provided.

Zootaxa 4237 (1): 078-090
<http://www.mapress.com/j/zt/>
 Copyright © 2017 Magnolia Press

Article

ISSN 1175-5326 (print edition)
ZOOTAXA
 ISSN 1175-5334 (online edition)

<https://doi.org/10.11646/zootaxa.4237.1.4>
<http://zoobank.org/urn:lsid:zoobank.org:pub:8E3AA8C7-B7B2-4AAE-BF12-D40138E5B61>

Biology and description of *Megaprosternum cleonarvorum* sp. nov. (Hymenoptera: Bethyidae) a gregarious larval ectoparasitoid of *Cleonaria bicolor* Thomson (Coleoptera: Cerambycidae) from India

ANKITA GUPTA^{1,2}, S. K. RAJESHWARI¹ & CELSO O. AZEVEDO³

¹ICAR-National Bureau of Agricultural Insect Resources, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bengaluru, 560 024, Karnataka, India. E-mail: shobini@gmail.com

²Universidade Federal do Espírito Santo, Departamento de Ciências Biológicas, Av. Fernando Ferrari 514, Goiabeiras, 29075-910, Vitória ES Brazil. E-mail: bethyidae@gmail.com

*Corresponding author. E-mail: drankitgupta7@gmail.com

Abstract

Megaprosternum Azevedo (Bethyidae: Scleroderminae) is newly recorded from the Oriental region, and *M. cleonarvorum* Gupta & Azevedo sp. nov. is described and illustrated from Southern India as a gregarious larval ectoparasitoid of *Cleonaria bicolor* Thomson (Coleoptera: Cerambycidae) on the host plant *Isora coccinea* L. (Rubiaceae). This is the first ever documentation of the biology of *Megaprosternum* Azevedo.

Zootaxa 4158 (2): 281-291
<http://www.mapress.com/j/zt/>
 Copyright © 2016 Magnolia Press

Article

ISSN 1175-5326 (print edition)
ZOOTAXA
 ISSN 1175-5334 (online edition)

<https://doi.org/10.11646/zootaxa.4158.2.9>
<http://zoobank.org/urn:lsid:zoobank.org:pub:BD49ECFE-84B7-4AE2-A2B2-52AF88D327AC>

A new species of *Crinibracon* Quicke (Hymenoptera: Braconidae) parasitic on pupae of *Hasora chromus* (Cramer) (Lepidoptera: Hesperidae) from India

ANKITA GUPTA¹, CORNELIS VAN ACHTERBERG² & MALATHI CHITRALA¹

¹ICAR-National Bureau of Agricultural Insect Resources, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bengaluru, 560 024, India.

²Dept. Terrestrial Zoology, Naturalis Biodiversity Center, Darwinweg 2, 2333 CR Leiden, Netherlands

*Corresponding author. E-mail: drankitgupta7@gmail.com

Abstract

A new species, *Crinibracon chromusae* Gupta & van Achterberg sp. n., parasitic on pupae of *Hasora chromus* (Cramer) (Hesperidae) on *Millettia* (=Pongamia) pinnata (L.) Panigrahi (Fabaceae), is described from India and compared with *C. sinicus* (Yang, Chen & Liu, 2008) from China, the only other species known with a similar general appearance. For the first time biological information for the genus *Crinibracon* Quicke, 1988, is given. Three species of hyperparasitoids, *Phaenolema braconidis* (Ferrière) (Hymenoptera: Eurytomidae), *Nesolynx javanica* Ferrière (Hymenoptera: Eulophidae), and an *Exepelus* sp. (Hymenoptera: Eupelmidae) emerged along with *C. chromusae* sp. n. from pupae of *H. chromus*. The generic placement of this new species along with interesting parasitoid biology is discussed.

Zootaxa 3827 (4): 439-470
www.mapress.com/zootaxa
 Copyright © 2014 Magnolia Press

Article

ISSN 1175-5326 (print edition)
ZOOTAXA
 ISSN 1175-5334 (online edition)

<http://dx.doi.org/10.11646/zootaxa.3827.4.2>
<http://zoobank.org/urn:lsid:zoobank.org:pub:F341F721-122C-41D8A-9B35-1FBFD166D5C0>

Lycanidae parasitoids from peninsular India with description of four new species of microgastrine wasps (Hymenoptera: Braconidae) along with new insights on host relationships

ANKITA GUPTA^{1,2}, PARESH V. CHURI¹, ASHOK SENGUPTA¹ & SARANG MHATRE¹

¹National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bengaluru, Karnataka, India, 56002

²Ram Mandir Trust Bldg, 14/B, 3rd Floor, Sitata Devi Temple Road, Mahim, Mumbai, 400016

³Kendriya Vidyalaya No. 1, Jalahalli (West), Bengaluru, 560015

⁴204, C wing, Gangadhar complex, Kamathe, Navi, Mumbai, 410209

*Corresponding author. E-mail: drankitgupta7@gmail.com

Abstract

In the comprehensive rearing of lepidopterans from peninsular India, eleven species of Lycanidae were parasitized by ten species of wasps. Four new taxa of lycanid associated microgastrine wasps *Parapanteles eros* Gupta n. sp., *P. arka* Gupta n. sp., *P. esha* Gupta n. sp., and *P. regale* Gupta n. sp. reared from *Chilades pandava* (Horsfield), *Caretta thersites* (Drury), *Proxostes dubiosa* (Semper), *Tajuria cippus* (Fabricius), respectively, are described with diagnostic details and illustrations along with documentation of six species of wasps viz., *Apanteles folia*, *Apanteles* sp., *Protopanteles* sp. 01 & 02 (Braconidae), *Choragob obtusus* Morley (Ichneumonidae), and *Brachymeria lasus* (Walker) (Chalcididae). This is the first record of host-parasitoid association of lycanid butterflies with *Parapanteles*. Wasps from three different families were recorded: Braconidae, Ichneumonidae, and Chalcididae. The parasitoid species were reared from the following Lycanidae hosts: *Anthrene lycanina* (Velder), *Arhopala amara* Hewitson, *Chilades pandava* (Horsfield), *Caretta thersites* (Drury), *Janitades celene* (Cramer), *Proxostes dubiosa* (Semper), *Rathinda emor* (Fabricius), *Spindaris vulcanus* (Fabricius), *Tajuria cippus* (Fabricius), *Tarucus baltanicus* nigra Bethune-Baker, and *Tarucus callimera* Butler. All lycanids were collected from peninsular India, except *T. callimera* (central India). A comparative account of all newly described species is provided along with the detailed illustrated description and differences vis-à-vis its closely allied Indian species. Also a comprehensive table comprising parasitoid species, associated host, stage of parasitism, and nature of cocoon is provided.

A new species of *Parapanteles* Ashmead (Hymenoptera: Braconidae) from India reared from *Abisara echeria* Stoll (Lepidoptera: Riodinidae) with key to the Indian *Parapanteles* species

ANKITA GUPTA^{1,4}, BLAISE PEREIRA¹ & PARESH V. CHURI¹

¹National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560 024, Karnataka, India

²Blaissette Plot 5, I. C. Colony, Cross Road No. 3, Borivli West, Mumbai 400103. E-mail: blaispereira@gmail.com

³Ram Mandir trust Bldg, 14/B, 3rd Floor, Sitla Devi Temple Road, Mahim, Mumbai-400016. E-mail: pareschuri@gmail.com

⁴Corresponding author. E-mail: drankitagupta7@gmail.com

Abstract

A new species of gregarious endoparasitoid, *Parapanteles echeriae* Gupta, Pereira & Churi, sp. nov., bred from *Abisara echeria* Stoll (Lepidoptera: Riodinidae) on host plant *Embelia* sp. (Myrsinaceae), is described and illustrated from Mumbai, Maharashtra, India. *Abisara echeria*, commonly known as plum jady, is a small striking butterfly prevalent in Asia. This is the first ever record of a parasitoid wasp associated with *Abisara*. *Parapanteles echeriae* is distinguished from *P. streptochaeta* Ahmad & Akhtar in having propodeum with areola 1.2x as long as wide; anterior diagonal carina meeting anterior margin of propodeum in one half; veins r and 2RS of fore wing merging in a smooth curve; yellow coloured legs and tegulae brownish black. A key to the Indian species of *Parapanteles* is also provided.

Parasitoids of Hesperidae from peninsular India with description of a new species of *Dolichogenidea* (Hymenoptera: Braconidae) parasitic on caterpillar of *Borbo cinnara* (Wallace) (Lepidoptera: Hesperidae)

ANKITA GUPTA¹, SWAPNIL A. LOKHANDE² & ABHAY SOMAN³

¹National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560024, Karnataka, India

²Dept. of Mathematics, IIT-Bombay, Powai, Mumbai-400076. E-mails: swapnilokhande@gmail.com, somanabhay@gmail.com

³Corresponding author E-mail: drankitagupta7@gmail.com

Abstract

Five species of parasitoid wasps associated with hesperids from peninsular India are documented along with the description of a new species of gregarious endoparasitoid, *Dolichogenidea cinnarae* sp. nov. (Hymenoptera: Braconidae) parasitic on caterpillar of *Borbo cinnara* (Wallace) (Lepidoptera: Hesperidae). Also, the gregarious larval parasitoid, *Cotesia erionotae* (Wilkinson) (Braconidae) and solitary pupal parasitoid *Chorops plusio* Gupta & Maheshwary (Ichneumonidae) were bred from the host *Udaspes folius* (Cramer) on the host plant *Hedyochium coronarium* J. Koenig. *Udaspes folius* is the new host record for the parasitoid wasp genus *Chorops*. *Cotesia erionotae* was bred from *U. folius* caterpillars from three states: Maharashtra, Karnataka and Kerala. An encyrtid wasp *Ooencyrtus papilionis* Ashmead was bred from eggs of *Bibasis jaita* (Moore) on the host plant *Hiptage benghalensis* (L.). This is the first documentation of a parasitoid wasp from the genus *Bibasis*. *Leptobotis indica* (Cameron) (Ichneumonidae), often associated with *Parnara guttatus* (Bremer & Grey), was recorded from the Andaman islands.

A new species of *Glyptapanteles* (Hymenoptera: Braconidae: Microgasterinae), a larval parasitoid of *Elymnias hypermnestra* (Linnaeus) (Lepidoptera: Nymphalidae), along with some new host records of parasitoids from Peninsular India

ANKITA GUPTA¹ & BLAISE PEREIRA²

¹National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560 024, Karnataka, India. E-mail: drankitagupta7@gmail.com

²Blaissette Plot 5, I. C. Colony, Cross Road No. 3, Borivli West, Mumbai 400103. E-mail: blaispereira@gmail.com

Abstract

A new species, *Glyptapanteles hypermnestrae* Gupta and Pereira, is described from Maharashtra, India, and compared with closely allied species. This new species was bred from parasitized larvae of *Elymnias hypermnestra* (Linnaeus) (Lepidoptera: Nymphalidae). In addition to this, two hymenopterian parasitoids, *Apanteles folia* Nixon (Braconidae: Microgasterinae) and *Brachymeria indica* (Krausse) (Chalcidoidea: Pteromalidae), are for first time reported parasitizing larvae of *Arhopala amantae* (Hewitson) (Lepidoptera: Lycaenidae) and pupae of *Parerionia valeria* (Cramer) (Lepidoptera: Pieridae) respectively.



A New Species of *Gonatocerus* Nees (Hymenoptera: Chalcidoidea: Mymaridae) from Karnataka, India

ANKITA GUPTA AND J. POORANI

Project Directorate of Biological Control (ICAR)
P. B. No. 2491, H. A. Farm post, Bellary Road, Hebbal
Bangalore 560 024, Karnataka, India

Corresponding author: ankitagupta9@rediffmail.com

ABSTRACT - *Gonatocerus similis* sp. nov. (Hymenoptera: Chalcidoidea: Mymaridae) belonging to the *asulcifrons*-group, from Karnataka, India, is described and illustrated. The diagnostic characters of *G. similis* in relation to the other species of the *asulcifrons*-group and species of other groups from India are discussed in brief: First funicular segments shorter than pedicel and second funicular segment; sixth and seventh funicular segments yellowish white; fore wing disc with 5 microtrichia; fore coxae yellowish; absence of subantennal sulci on head; antenna with funicular segments 2-8 subequal and clava with more than 11 sensillae.

Three new species of reared parasitic wasps (Hymenoptera: Braconidae: Microgasterinae) from India

ANKITA GUPTA

National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560 024, Karnataka, India. E-mail: drankitagupta7@gmail.com

Abstract

Three new species of parasitic wasps (Braconidae: Microgasterinae) from southern India are described and illustrated. *Glyptapanteles clanisae* sp. nov., a remarkable gregarious endoparasitoid, was bred from the caterpillar of *Clanis phalaris* Cramer (Lepidoptera: Spingidae) on the host plant *Pongamia pinnata* (L.) (Leguminosae) along with a hyperparasitoid, *Eurytoma* sp. (Eurytomidae). *Glyptapanteles trilochae* sp. nov., was reared from parasitized caterpillar of *Trilocha varians* (Walker) (Lepidoptera: Bombycidae) on the host plant *Ficus racemosa* L. (Moraceae) along with a hyperparasitoid, *Paraphylax* sp. (Ichneumonidae: Cryptinae). *Buluka horni* sp. nov. was collected from solitary cocoons of an indeterminate caterpillar feeding on *Mangifera indica* L. leaves. This study also confirms a host range extension of Indian species of *Glyptapanteles* to Bombycidae and Spingidae in addition to the earlier documented families viz., Papilionidae, Nymphalidae, Arctiidae, and Noctuidae.

Revision of the Indian *Microplitis* Foerster (Hymenoptera: Braconidae: Microgasterinae), with description of one new species

ANKITA GUPTA

National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560 024, Karnataka, India. E-mail: drankitagupta7@gmail.com

Abstract

The Indian fauna of the genus *Microplitis* Foerster, 1862 is revised. An illustrated key to eight species including the description of one new species, *M. markyi* sp. nov., is provided. Six previously described species, namely: *M. ajmerensis* Rao & Kurian, *M. demolitor* Wilkinson, *M. indicus* Marsh., *M. manilae* Ashmead, *M. prodeniae* Rao & Kurian, and *M. spodiopetiae* Rao & Kurian are elaborated with taxonomical variations and extended distribution. Two species, *M. baghesri* Sathe, Inamdar & Dawale and *M. dipika* (Bhatnagar) are considered *incertae sedis*. A new combination is suggested for *Snellenius maculipennis* (Szépligeti) which is placed into synonymy with *Microplitis*. Information on taxonomic history of the genus, diagnostic characters of all the included species, distribution and host relationships are provided.

Reared parasitic wasps attacking hesperids from Western Ghats (Kerala, India) with description of a new species of *Dolichogenidea* (Hymenoptera: Braconidae) as a larval parasitoid of *Thoreasa evershedi* (Evans) (Lepidoptera: Hesperidae)

ANKITA GUPTA¹ & KALESH S²

¹National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560 024, Karnataka, India

²Greeham, BN 439, Bapuji Nagar, Medical College Post, Thiruvananthapuram, Kerala, India. E-mail: kaleshs2002ie@gmail.com

³Corresponding author. E-mail: drankitagupta7@gmail.com

Abstract

Eight species of parasitic wasps were bred from various stages of five species of hesperids viz. *Thoreasa evershedi* (Evans), *Pelipotis mathias* (Fabricius), *Udaspes folius* (Cramer), *Borbo cinnara* (Wallace) and *Calotis* sp. inhabiting Western Ghats, Kerala, India. One new species, *Dolichogenidea kushi* Gupta & Kalesh, is described and illustrated from Kerala, India, and its relationship with closely allied species is discussed. This new species was bred from parasitized larvae of *Thoreasa evershedi* (Evans) (Lepidoptera: Hesperidae). Microgasterinae braconid species, *Apanteles juvenis* Rohwer and *Cotesia erionotae* (Wilkinson) were bred from parasitized larvae of *P. mathias* and *U. folius* respectively. *Brachymeria kushi* Oudikmen (Chalcidoidea) was recorded from pupae of *U. folius*. Two hyperparasitoids, *Eurytoma manilae* Ashmead (Eurytomidae) & *Pedobius foveolatus* (Crawford) (Aphelinidae) were bred from larvae of *B. cinnara*. Egg parasitoid, *Ooencyrtus papilionis* Ashmead (Encyrtidae) and pupal parasitoid *Brachymeria laus* (Walker) (Chalcidoidea) was recorded from *Calotis* sp. Information on the parasitoid distribution, brief diagnosis of each species with a habitus photograph for easy identification, host association, host caterpillars, caterpillar host plants, and taxonomic comments are provided.

A NEW SPECIES OF *NEASTYMACHUS* GIRAUT (HYMENOPTERA: ENCYRTIDAE) FROM KARNATAKA STATE OF INDIA

MOHAMMAD HAYAT^{1,3}, ANKITA GUPTA² & KRISHNAN SELVARAJ²

¹Department of Zoology, Aligarh Muslim University, Aligarh 202 002, India.

²ICAR-National Bureau of Agricultural Insect Resources, Bengaluru 560 024, India.

³Corresponding author. E-mail: havat.mohd44@gmail.com

Abstract. A new species of the genus *Neastymachus* Girault (Hymenoptera: Encyrtidae), *N. notialis* Hayat & Gupta, sp. nov., is described from a specimen collected in Udipi, Karnataka (India). This species is compared with two apparently similar species, *N. angustifrons* (Shafee et al., 1975) and *N. katekenis* Hayat, in Hayat et al. (2015).

Keywords: Encyrtinae, Discodini, new species, India.

http://dx.doi.org/10.11646/zootaxa.3616.6.1
http://zoobank.org/urn:lsid:zoobank.org:pub:E1F586D6-28D6-445A-986F-5C08839E834E

Microgastrinae (Hymenoptera: Braconidae) of Reunion Island: a catalogue of the local species, including 18 new taxa and a key to species

PASCAL ROUSSE¹ & ANKITA GUPTA²

¹ Corresponding author: Iziko South African Museum, Natural History Department, PO Box 61, Cape Town 8000, Sth Africa / Stellenbosch University, Department of Botany and Zoology, Evolutionary Genomics Group, Private Bag X1, Stellenbosch 7602, Sth Africa.
rousse.pascal@wanadoo.fr

² National Bureau of Agriculturally Important Insects, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560 024, Karnataka, India.
drankitagupta7@gmail.com

Abstract

Microgastrine wasps (Hymenoptera: Braconidae) are primary parasitoids of Lepidoptera. Some of them parasitise major economic pests and have been largely used for biocontrol programs. We revise here the fauna of Reunion Island: 34 species were recorded, belonging to 13 genera. One genus and 18 species are newly described: *Dodogaster* gen. nov., *Apanteles minatchy* sp. nov., *A. pashmina* sp. nov., *A. romei* sp. nov., *Cotesia xavieri* sp. nov., *Dolichogenidea ashoka* sp. nov., *D. broadi* sp. nov., *D. lumba* sp. nov., *D. uru* sp. nov., *D. villemantae* sp. nov., *Distatrix yunae* sp. nov., *Dodogaster grangeri* sp. nov., *Exoryza saffranum* sp. nov., *Glyptapanteles chidra* sp. nov., *Nyereria ganges* sp. nov., *N. mayurus* sp. nov., *Parapanteles covino* sp. nov., *P. dagnac* sp. nov. and *Wilkinsonellus narangahus* sp. nov.. On the remaining 16 species, 12 are new distribution records. Finally, *Glyptapanteles antsirabensis* (Granger) comb. nov., *G. ficus* (Granger) comb. nov., *G. subandinus* (Blanchard) comb. nov., and *Venandides curticornis* (Granger) comb. nov. are reassigned from the traditionally defined *Apanteles* genus to these respective genera. A key to all genera and species is provided.

J. Biol. Control, 21(2): 287-289, 2007

Research Note



J. Biol. Control, 21(2): 291-293, 2007

Research Note



A new species of *Trichogramma* (Hymenoptera: Trichogrammatidae) from Anantnag, Jammu & Kashmir, India

H. NAGARAJA¹*, JAMAL AHMAD² and ANKITA GUPTA¹

¹Project Directorate of Biological Control, Post Bag No. 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore – 560024, Karnataka, India

²Division of Entomology, Sher-e-Kashmir University of Agricultural Sciences and Technology (K), Shalimar Campus, Srinagar – 191121, Jammu & Kashmir, India

*E-mail: hulikalnagaraja@yahoo.com

A new species of *Trichogramma* (Hymenoptera: Trichogrammatidae) parasitic on eggs of pomegranate fruit borer, *Deudorix isocrates* (Fabricius) (Lepidoptera: Lycaenidae)

H. NAGARAJA and ANKITA GUPTA

Project Directorate of Biological Control, Post Bag No. 2491, H.A. Farm Post, Bellary Road, Hebbal, Bangalore – 560024, Karnataka, India

E-mail: hulikalnagaraja@yahoo.com

ENTOMON 32(3): 161–167 (2007)
Article No. ent.32302

ABSTRACT:



new egg parasitoid of pomegranate fruit Lycaenidae), is described and illustrated

A new species of *Dinarmus* Thomson (Hymenoptera: Chalcidoidea: Pteromalidae) from India

Ankita Gupta

Project Directorate of Biological Control, P. B. No 2491, H. A. Farm Post, Bellary Road, Hebbal, Bangalore 560 024, India
Email: ankitagupta9@rediffmail.com

ABSTRACT: *Dinarmus crotalariae* sp. nov. is described from Karnataka, India and a modified key to the Indian species of *Dinarmus* given.

© 2007 Association for Advancement of Entomology